

Spin stiffness in the frustrated Heisenberg antiferromagnet

M.S.L. du Croo de Jongh, P.J.H. Denteneer

Instituut-Lorentz, University of Leiden, P. O. Box 9506, 2300 RA Leiden, The Netherlands

Abstract

We calculate the spin stiffness of the $S = \frac{1}{2}$ frustrated Heisenberg antiferromagnet directly from a general formula which is evaluated in the Schwinger boson mean-field approximation. Both Néel and collinear ordering are considered. For collinear ordering, we take the anisotropy of this phase into account, unlike previous approaches. For Néel ordering, a detailed study is made of the finite-size scaling behavior of the two terms that make up the spin stiffness. The exponents of the scaling with the system size of the two terms comprising the spin stiffness turn out to be identical to those of the unfrustrated case.

PACS numbers : 75.10.Jm, 75.30.Kz, 75.40.Cx

I. INTRODUCTION

The recent interest in the frustrated Heisenberg antiferromagnets is motivated by high T_c -superconductivity; the undoped compounds show long-range antiferromagnetic order, similar to the Heisenberg model. Upon doping superconductivity occurs. Adding frustration to the Heisenberg model can be thought of as to mimic the effect of hole doping.

We consider the frustrated Heisenberg model on a square lattice with $N = L^2$ sites. It is described by the following Hamiltonian for quantum spins $\mathbf{S}_j$ on a lattice:

$$H = J_1 \sum_{nn} \mathbf{S}_i \cdot \mathbf{S}_j + J_2 \sum_{nnn} \mathbf{S}_i \cdot \mathbf{S}_j, \quad (1)$$

where nn denotes a pair (ij) of nearest-neighbor sites and nnn a pair of next-nearest-neighbor sites. The spin length is fixed; $S = \frac{1}{2}$. Both J_1 and J_2 are taken to be non-negative. If

J_2/J_1 is small, the antiferromagnetic long-range order is recovered (Néel-like). For J_2/J_1 large, the system decomposes in two Néel ordered sublattices which, however, have the same quantization axis. Alternating strips of up and down-spins will occur; the so-called collinear ordering. Clearly these couplings frustrate each other. If the spins were classical objects a large number of phases, among which the Néel and collinear phases, would become degenerate for $J_2/J_1 = 0.5$. For the quantum case, a quantum phase transition to a spin-liquid phase might occur.¹⁻⁹

In this paper, our intention is to employ the spin stiffness ρ_s to measure magnetic order in the system. We calculate the spin stiffness in the framework of the Schwinger boson mean-field approximation (SBMFA) using a general formula for ρ_s . Previous evaluations of the spin stiffness were indirect^{3,11,12}. However, apart from minor adjustments, we confirm their results. Furthermore the scaling behavior of ρ_s in this approximation is derived. This is useful to sensibly extrapolate results of more exact approaches, like quantum Monte Carlo and exact diagonalization. Also, the typical system sizes for which scaling is valid can be estimated in this way.⁵⁻⁷

II. SCHWINGER BOSON MEAN-FIELD APPROXIMATION

The SBMFA improves upon standard mean-field theory by incorporating correlations between neighboring spins^{10,11}. With this approximation we derive the energies and wavefunctions of all states of the frustrated Heisenberg model for both Néel and collinear order. Our notation below will generally follow that of Mila et al.²

The Schwinger boson transformation is a representation of the separate spin operators by pairs of boson operators; $S^+ = a^\dagger b$, $S^- = ab^\dagger$, $S^z = \frac{1}{2} [a^\dagger a - b^\dagger b]$, supplemented by the local constraint $a^\dagger a + b^\dagger b = 2S(=1)$.

To transform the Hamiltonian in a convenient form an appropriate rotation in spin space is applied. Define $D_{ij} = a_i a_j^\dagger + b_i b_j^\dagger$ and $B_{ij} = a_i b_j + b_i a_j$. The Hamiltonian becomes

$$H = -\frac{1}{2} \sum_{AFM} J_{ij} (B_{ij}^\dagger B_{ij} - \frac{1}{2}) + \frac{1}{2} \sum_{FM} J_{ij} (D_{ij}^\dagger D_{ij} - \frac{3}{2}), \quad (2)$$

where we have inserted D_{ij} for the pairs (ij) of spins parallel in the S^z -direction and B_{ij} for the antiparallel pairs. For the two orderings considered this is depicted in Figure 1. The parameter J_{ij} equals J_1 for nearest-neighbors and J_2 for next-nearest-neighbors. The mean-field decoupling is made using the fields $\kappa_{ij} = \frac{1}{2}\langle D_{ij} \rangle$ and $\gamma_{ij} = \frac{1}{2}\langle B_{ij} \rangle$. The local constraint $a^\dagger a + b^\dagger b = 1$ is replaced by a global one and enforced by means of a Lagrange multiplier λ . After a Fourier transform from (a_i, b_i) to $(a_{\mathbf{p}}, b_{\mathbf{p}})$ and a Bogoliubov transformation from $(a_{\mathbf{p}}, b_{\mathbf{p}})$ to $(\alpha_{\mathbf{p}}, \beta_{\mathbf{p}})$;

$$\begin{aligned} a_{\mathbf{p}} &= \alpha_{\mathbf{p}} \cosh \theta_{\mathbf{p}} + \beta_{\mathbf{p}}^\dagger \sinh \theta_{\mathbf{p}}, \\ b_{-\mathbf{p}}^\dagger &= \alpha_{\mathbf{p}} \sinh \theta_{\mathbf{p}} + \beta_{\mathbf{p}}^\dagger \cosh \theta_{\mathbf{p}}, \\ \tanh 2\theta_{\mathbf{p}} &= \frac{\Delta_{\mathbf{p}}}{h_{\mathbf{p}} + \lambda}, \end{aligned} \quad (3)$$

the Hamiltonian becomes

$$H_{MF} = E_c + \sum_{\mathbf{p}} \omega_{\mathbf{p}} (\alpha_{\mathbf{p}}^\dagger \alpha_{\mathbf{p}} + \beta_{\mathbf{p}} \beta_{\mathbf{p}}^\dagger), \quad (4)$$

where we have introduced the quasiparticle energies

$$\omega_{\mathbf{p}} = \sqrt{(h_{\mathbf{p}} + \lambda)^2 - \Delta_{\mathbf{p}}^2}. \quad (5)$$

Below the quantities $h_{\mathbf{p}}$, $\Delta_{\mathbf{p}}$ and E_c will be defined for the Néel and collinear orderings separately. The fields κ_{ij} , γ_{ij} and the Lagrange multiplier λ are obtained through consistency equations.

We consider two types of order:

Néel order: the specific form of $h_{\mathbf{p}}$, $\Delta_{\mathbf{p}}$, E_c and the three consistency equations are:

$$h_{\mathbf{p}} = 4J_2\kappa \cos p_x \cos p_y, \quad (6)$$

$$\Delta_{\mathbf{p}} = 2J_1\gamma(\cos p_x + \cos p_y), \quad (7)$$

$$E_c = 2N \left[J_1 \left(\frac{1}{4} + 2\gamma^2 \right) - J_2 \left(\frac{3}{4} + 2\kappa^2 \right) - \lambda \right], \quad (8)$$

$$\kappa = \frac{1}{N} \sum_{\mathbf{p}} \frac{h_{\mathbf{p}} + \lambda}{2\omega_{\mathbf{p}}} \cos p_x \cos p_y, \quad (9)$$

$$\gamma = \frac{1}{N} \sum_{\mathbf{p}} \frac{\Delta_{\mathbf{p}}}{4\omega_{\mathbf{p}}} (\cos p_x + \cos p_y), \quad (10)$$

$$1 = \frac{1}{N} \sum_{\mathbf{p}} \frac{h_{\mathbf{p}} + \lambda}{2\omega_{\mathbf{p}}}. \quad (11)$$

For future applications it is also useful to define the “condensate” m_s ¹³ by

$$m_s = \frac{h_{\mathbf{0}} + \lambda}{N\omega_{\mathbf{0}}} = 1 - \frac{1}{N} \sum_{\mathbf{p} \neq (0,0), (\pi,\pi)} \frac{h_{\mathbf{p}} + \lambda}{2\omega_{\mathbf{p}}} \quad (12)$$

This is the combination of the -equivalent- $\mathbf{p} = (0,0)$ and $\mathbf{p} = (\pi,\pi)$ terms in (9) and (11), which both diverge for $N \rightarrow \infty$. In the same limit, $N \rightarrow \infty$, m_s also equals the corresponding terms in (10).

Collinear order: for this phase we introduce quantities with a bar where confusion might arise;

$$\bar{h}_{\mathbf{p}} = 2J_1 \bar{\kappa} \cos p_x,$$

$$\bar{\Delta}_{\mathbf{p}} = 2J_1 \gamma_1 \cos p_y + 4J_2 \gamma_2 \cos p_x \cos p_y, \quad (13)$$

$$\bar{E}_c = 2N \left(J_1 (\gamma_1^2 - \bar{\kappa}^2 - \tfrac{1}{4}) + J_2 (\tfrac{1}{4} + 2\gamma_2^2) - \bar{\lambda} \right), \quad (14)$$

$$\bar{\kappa} = \frac{1}{N} \sum_{\mathbf{p}} \frac{\bar{h}_{\mathbf{p}} + \bar{\lambda}}{2\bar{\omega}_{\mathbf{p}}} \cos p_x, \quad (15)$$

$$\gamma_1 = \frac{1}{N} \sum_{\mathbf{p}} \frac{\bar{\Delta}_{\mathbf{p}}}{2\bar{\omega}_{\mathbf{p}}} \cos p_y, \quad (16)$$

$$\gamma_2 = \frac{1}{N} \sum_{\mathbf{p}} \frac{\bar{\Delta}_{\mathbf{p}}}{2\bar{\omega}_{\mathbf{p}}} \cos p_x \cos p_y, \quad (17)$$

$$1 = \frac{1}{N} \sum_{\mathbf{p}} \frac{\bar{h}_{\mathbf{p}} + \bar{\lambda}}{2\bar{\omega}_{\mathbf{p}}}. \quad (18)$$

The condensate $\bar{m}_s$ and $\bar{\omega}_{\mathbf{p}}$ are defined in a similar manner as before in (12) and (5).

The discussion above has given us the ground states $|0\rangle$ with energy $E_0 = E_c + \sum_{\mathbf{p}} \omega_{\mathbf{p}}$ for both orderings. These ground states are characterized by the absence of quasi-particles; $\alpha|0\rangle = \beta|0\rangle = 0$. Excited states are given by:

$$|a\rangle = \alpha_1^\dagger \cdot \dots \cdot \alpha_n^\dagger \cdot \beta_1^\dagger \cdot \dots \cdot \beta_m^\dagger |0\rangle, \quad (19)$$

$$\begin{aligned}
H_{MF}|a\rangle &= E_a|a\rangle \\
&= [E_0 + \omega_1 + \dots + \omega_n + \omega_1 + \dots + \omega_m]|a\rangle.
\end{aligned} \tag{20}$$

III. THE SPIN STIFFNESS IN THE SBMFA

The spin stiffness ρ_s is non-zero if there exists magnetic order in the system and is, at $T = 0$, associated with an increase in energy upon twisting the order parameter of the system ($\Delta E = \frac{1}{2}N\rho_s|\mathbf{q}|^2$ with $\mathbf{q}$ the wave vector of the twist.)^{15,14}. In line with Einarsson and Schulz⁵ we introduce this twist through a replacement of D_{ij} and B_{ij} in (2) by

$$D_{ij}(\mathbf{q}) = a_i a_j^\dagger e^{\frac{i}{2}\mathbf{q}\cdot\mathbf{r}_\delta} + b_i b_j^\dagger e^{-\frac{i}{2}\mathbf{q}\cdot\mathbf{r}_\delta} \tag{21}$$

$$B_{ij}(\mathbf{q}) = a_i b_j e^{\frac{i}{2}\mathbf{q}\cdot\mathbf{r}_\delta} + b_i a_j e^{-\frac{i}{2}\mathbf{q}\cdot\mathbf{r}_\delta}, \tag{22}$$

with $\mathbf{r}_\delta = \mathbf{r}_j - \mathbf{r}_i$. The resulting Hamiltonian $H(\mathbf{q})$ is now evaluated within the SBMFA.

Defining $\kappa_{ij}(\mathbf{q}) = \frac{1}{2}\langle D_{ij}(\mathbf{q}) \rangle$ and $\gamma_{ij}(\mathbf{q}) = \frac{1}{2}\langle B_{ij}(\mathbf{q}) \rangle$, the mean-field Hamiltonian becomes

$$\begin{aligned}
H_{MF}(\mathbf{q}) &= - \sum_{AFM} J_{ij} \gamma_{ij}(\mathbf{q}) (B_{ij}^\dagger(\mathbf{q}) + B_{ij}(\mathbf{q}) - 2\gamma_{ij}(\mathbf{q})) \\
&\quad + \sum_{FM} J_{ij} \kappa_{ij}(\mathbf{q}) (D_{ij}^\dagger(\mathbf{q}) + D_{ij}(\mathbf{q}) - 2\kappa_{ij}(\mathbf{q})) \\
&\quad + \lambda \sum_i (a_i^\dagger a_i + b_i^\dagger b_i - 1) + \text{constants}.
\end{aligned} \tag{23}$$

Since within the SBMFA we know what the excited states are [see (19) and (20)], the spin stiffness can be directly evaluated from second order perturbation theory

$$\rho_s = -\frac{1}{N}\langle 0|t|0\rangle + \frac{2}{N} \sum_{|a\rangle \neq |0\rangle} \frac{|\langle 0|j|a\rangle|^2}{E_0 - E_a} \equiv T + J, \tag{24}$$

with the quantities t and j defined by

$$t = -\frac{d^2}{dq^2} H_{MF}(\mathbf{q}) \Big|_{\mathbf{q}=0}, \quad j = \frac{d}{dq} H_{MF}(\mathbf{q}) \Big|_{\mathbf{q}=0}. \tag{25}$$

In (24), we have also defined the abbreviations T and J for the two terms in ρ_s .

IV. RESULTS FOR THE SPIN STIFFNESS

From this point on, we set $\mathbf{q} = q(\cos \phi, \sin \phi)$. Of the two terms for ρ_s in (24), T is evaluated more easily. We obtain after straightforward manipulations with Brillouin zone summations :

$$T = J_1 \gamma^2 - 2J_2 \kappa^2, \quad (26)$$

$$\bar{T} = 2J_2 \gamma_2^2 + J_1 (\gamma_1^2 \sin^2 \phi - \bar{\kappa}^2 \cos^2 \phi), \quad (27)$$

These simple equations hold for all system sizes N .

The quantity J requires more effort; the operator j has to be expressed in the operators $\alpha_{\mathbf{p}}$ and $\beta_{\mathbf{p}}$ defined in (3). For the wavefunctions $|a\rangle$ and energies E_a of the excitations we use (19) and (20). The resulting values for J are written as summations over the Brillouin zone:

$$\begin{aligned} J &= -\frac{1}{N} \sum_{\mathbf{p}} \frac{\sin^2 p_y}{\omega_{\mathbf{p}}^3} (J_1 \gamma (h_{\mathbf{p}} + \lambda) - 2J_2 \kappa \Delta_{\mathbf{p}} \cos p_x)^2, \\ \bar{J} &= -\frac{1}{N} \sum_{\mathbf{p}} \frac{1}{\bar{\omega}_{\mathbf{p}}^3} (\cos \phi \sin p_x [J_1 \bar{\kappa} \bar{\Delta}_{\mathbf{p}} - 2J_2 \gamma_2 (\bar{h}_{\mathbf{p}} + \bar{\lambda}) \cos p_y] \\ &\quad - \sin \phi \sin p_y [J_1 \gamma_1 (\bar{h}_{\mathbf{p}} + \bar{\lambda}) + 2J_2 \gamma_2 (\bar{h}_{\mathbf{p}} + \bar{\lambda}) \cos p_x])^2. \end{aligned}$$

Only for the infinitely large lattice these equations can be simplified by replacing summations by integrals and partially integrating. The expression for the spin stiffness $\rho_s = T + J$ then simplifies considerably and becomes:

$$\rho_s = m_s (J_1 \gamma - 2J_2 \kappa), \quad (28)$$

$$\begin{aligned} \bar{\rho}_s &= \bar{m}_s [(2J_2 \gamma_2 - J_1 \bar{\kappa}) \cos^2 \phi \\ &\quad + (2J_2 \gamma_2 + J_1 \gamma_1) \sin^2 \phi]. \end{aligned} \quad (29)$$

Ivanov and Ivanov³ apply a different method to derive ρ_s . They use a modified spin wave theory which leads to the same consistency equations as the Schwinger boson approach. ρ_s is then obtained by calculating the correlation length ξ associated with the spin-spin correlation function $\langle \mathbf{S}_i \cdot \mathbf{S}_j \rangle$ and comparing this ξ to the expression for ξ obtained for the

non-linear sigma model to two-loop order by Chakravarty et al.¹² (where $\xi \sim \exp(2\pi\rho_s/\theta)$ with θ the temperature). For Néel ordering their expression is identical to our result (28). It is gratifying to see that the non-linear sigma model also is the effective field theory for the low-energy physics of the *frustrated* Heisenberg antiferromagnet.

For the collinear ordering they obtain the geometrical average of our $\cos^2 \phi$ and $\sin^2 \phi$ -terms, whereas we take explicitly the anisotropy of this phase into account. Still both expressions for $\bar{\rho}_s$ vanish at the same value of J_2/J_1 .

Table I and Figure 1 contain our numerical results.

V. SCALING OF THE SPIN STIFFNESS

It is necessary to know the size dependence of observables to obtain a good approximation for their limit values. Neuberger and Ziman¹⁶ derived the scaling behavior for an unfrustrated Heisenberg antiferromagnet explicitly. Here we extend this to the case of frustration.

Recently some discussion has arisen about where the scaling behavior of ρ_s sets in⁶. Our formulas in the last section lend themselves well to investigate this.

Here we only treat the Néel ordering. We want to know the scaling behavior of the condensate m_s and the two terms J and T that make up ρ_s ($\rho_s = T + J$). The latter two will turn out to have different scaling behavior.

As can be seen from (26) only the scaling behavior of κ_N and γ_N is required for T . These two are part of the set $(\kappa_N, \gamma_N, \lambda_N)$ of mutually dependent quantities. We will now argue what is the exponent of their scaling behavior and therefore of T , without trying to obtain the precise prefactor (which would be quite tedious).

Name the equations (9)-(11) *I*, *II*, and *III*, respectively. They contain poles at $\mathbf{p} = (0, 0)$ and $\mathbf{p} = (\pi, \pi)$. With help of (5), (6) and (7) we rearrange them as $I - III$, $II - III$ and $4J_2\kappa_N I - 4J_1\gamma_N II + \lambda_N III$. We neglect the $\mathbf{p} = (0, 0)$ and $\mathbf{p} = (\pi, \pi)$ terms in the summations. It is easy to show that this will give rise to errors of the order $O(N^{-2})$. Next

we expand these equations to first order round their infinite size values $(\kappa, \gamma, \lambda)$. Define the size dependences $\delta\kappa_N = \kappa_N - \kappa$, $\delta\gamma_N = \gamma_N - \gamma$ and $\delta\lambda_N = \lambda_N - \lambda$ to obtain the equation

$$\begin{bmatrix} \kappa - 1 \\ \gamma - 1 \\ \lambda - 4J_1\gamma^2 + 4J_2\kappa^2 \end{bmatrix} = \vec{A} + \vec{B} \cdot \begin{bmatrix} \delta\kappa_N \\ \delta\gamma_N \\ \delta\lambda_N \end{bmatrix}, \quad (30)$$

where $\vec{A}$ and $\vec{B}$ contain summations over the Brillouin zone dependent on the infinite size parameters κ , γ , and λ . The remaining size dependence of $\vec{B}$ can be neglected as it leads to higher order terms. On the other hand the summations in $\vec{A}$ will be replaced by integrations plus size dependent corrections. Using Neuberger and Ziman¹⁶ we obtain

$$\vec{B} \cdot \begin{bmatrix} \delta\kappa_N \\ \delta\gamma_N \\ \delta\lambda_N \end{bmatrix} = \frac{1}{N^{3/2}} \vec{C}. \quad (31)$$

The parameters κ_N , γ_N and λ_N thus scale with $N^{-3/2}$. A direct consequence of this is that $T_N - T \sim N^{-3/2}$. If the size dependence of the parameters is neglected $[(\kappa_N, \gamma_N, \lambda_N) \rightarrow (\kappa, \gamma, \lambda) \text{ inside the summations (9)-(11)}]$, we also find the prefactor:

$$T_N - T = \frac{0.7186}{N^{3/2}} \frac{1}{\sqrt{2}} \sqrt{\frac{\lambda + 4J_2\kappa}{\lambda - 4J_2\kappa}} [\lambda - 8J_2\kappa]. \quad (32)$$

Upto $J_2/J_1 \approx 0.5$ this is in good agreement with the numerical solution of (9)-(11) for various sizes N .

Next we consider the condensate $m_{s,N}$ and J_N , for which we can even derive the prefactors. Replacement of $(\kappa_N, \gamma_N, \lambda_N)$ by $(\kappa, \gamma, \lambda)$ in the summation of (12) leads to errors of the order $O(N^{-3/2})$. We will neglect these. Neuberger and Ziman¹⁶ state a lemma applicable to this summation. This leads to:

$$m_{s,N} - m_s = \frac{0.6208}{\sqrt{N}} \sqrt{2} \sqrt{\frac{\lambda + 4J_2\kappa}{\lambda - 4J_2\kappa}}. \quad (33)$$

For J_N we find in a similar fashion:

$$J_N - J = \frac{0.6208}{\sqrt{N}} \frac{\sqrt{\lambda^2 - (4J_2\kappa)^2}}{4\sqrt{2}}. \quad (34)$$

These formulas are in excellent agreement with the numerical solution to (9)-(11). For J_N this is depicted in figure 2.

In conclusion, we have obtained the scaling behavior of T and J from an analysis of the formulas in the SBMFA for the frustrated Heisenberg antiferromagnet. The qualitative scaling behavior (i.e. the exponents) is the same as for the *unfrustrated* case, which was discussed by Neuberger and Ziman¹⁶. The scaling behavior that was utilized by Einarsson and Schulz⁵ does agree with our findings but we confirm the message of Feiguin et al.⁶ that the clusters they used are too small for the scaling behavior of J to have set in. Numerically we see that the scaling behavior starts around size $N = 100$ (See Figure 2) whereas the largest cluster they used is $N = 36$.

We would like to thank J.M.J. van Leeuwen and W. van Saarloos for fruitful discussions.

REFERENCES

- ¹ P. Chandra, B. Doucot, Phys. Rev. B **38**, 9335 (1988).
- ² F. Mila, D. Poilblanc, C. Bruder, Phys. Rev. B **43**, 7891 (1991).
- ³ N.B. Ivanov, P. Ch. Ivanov, Phys. Rev. B **46**, 8206 (1992).
- ⁴ A.V. Dotsenko, O.P. Sushkov, Phys. Rev. B **50**, 13821 (1994).
- ⁵ T. Einarsson, H.J. Schulz, Phys. Rev. B **51**, 6151 (1995).
- ⁶ A.E. Feiguin, G.J. Gazza, A.E. Trumper, H.A. Ceccatto, Phys. Rev. B **52**, 15043 (1995).
- ⁷ H.J. Schulz, T.A.L. Ziman, D. Poilblanc, J. Phys. 1 France **6**, 675 (1996).
- ⁸ O.P. Sushkov, cond-mat 9602150.
- ⁹ M. Zhitomirsky, K. Ueda, Phys. Rev. B **54**, 9007 (1996)
- ¹⁰ D.P. Arovas, A. Auerbach, Phys. Rev. B **38**, 316 (1988), A. Auerbach, D.P. Arovas, Phys. Rev. Lett. **61**, 617 (1988).
- ¹¹ A. Auerbach, *Interacting Electrons and Quantum Magnetism* (Springer, New York, 1994).
- ¹² S. Chakravarty, B.I. Halperin, D.R. Nelson, Phys. Rev. B **39**, 2344 (1988).
- ¹³ J.E. Hirsch, S. Tang, Phys. Rev. B **39**, 2850 (1989).
- ¹⁴ M.E. Fisher, M.N. Barber, D. Jasnow, Phys. Rev. A **188**, 1111 (1973).
- ¹⁵ J.M.J. van Leeuwen, M.S.L. du Croo de Jongh, P.J.H. Denteneer, J. Phys. A:Math. Gen. **29**, 41 (1996).
- ¹⁶ H. Neuberger, T. Ziman, Phys. Rev. B **39**, 2608 (1989).

TABLES

J_2/J_1	Order	T		$-J$		ρ_s	
0.0	N	0.3352		0.1596		0.1757	
0.1	N	0.2961		0.1596		0.1365	
0.2	N	0.2597		0.1597		0.1000	
0.3	N	0.2271		0.1600		0.0672	
0.4	N	0.1995		0.1604		0.0391	
0.5	N	0.1783		0.1612		0.0171	
0.6	N	0.1639		0.1622		0.0017	
0.6	C	0.6231	0.0499	0.3841	0.0284	0.2390	0.0214
0.7	C	0.7211	0.1563	0.3327	0.0797	0.3884	0.0766
0.8	C	0.7794	0.2566	0.3232	0.1264	0.4563	0.1302
0.9	C	0.8327	0.3492	0.3292	0.1689	0.5034	0.1803
1.0	C	0.8866	0.4361	0.3434	0.2086	0.5432	0.2275
1.1	C	0.9420	0.5190	0.3623	0.2465	0.5797	0.2725

TABLE I. The limit ($N = \infty$) values for T , $-J$ and ρ_s as function of the ratio J_2/J_1 where $J_1 = 1$. The two orderings considered are Néel (N) and collinear (C) order. For the collinear ordering there are two directions: along the antiparallel spins, $\phi = \pi/2$ (listed first) and along the parallel spins, $\phi = 0$ (listed secondly).

FIGURES




FIG. 1. The spin stiffness ρ_s in units of J_1 (solid lines). For the collinear ordering the spin stiffness in the direction of the parallel spins, $\phi = 0$, (lower solid curve) and in the direction of the antiparallel spins $\phi = \pi/2$ (upper solid curve) are drawn. The dotted line is the result found by Ivanov and Ivanov³. *Inset* : The mean fields for the Néel (a) and the collinear order (b). Mind $\kappa \sim D_{ij}$ and $\gamma \sim B_{ij}$.




FIG. 2. The numerical scaling behavior of J_N (solid lines) for sizes $L = 2, 4, 6, 8, 10, 20, 40, 100$ (numbering is top-down) compared with the theoretical formula (34) (dotted line).